

IMAGES IN INTERVENTION

Case of Percutaneous Extracorporeal Femoro-Femoral Bypass for Acute Limb Ischemia From Large Bore Access




Saurav Chatterjee, MD, Riyaz Bashir, MD, RVT, Vladimir Lakhter, DO, Brian O'Murchu, MD, Brian O'Neill, MD, Vikas Aggarwal, MD, MPH, RPVI

An 80-year-old man presented with acute inferior ST-segment elevation myocardial infarction and cardiogenic shock. Cardiac catheterization revealed 99% stenosis in the mid-right coronary artery, a 95% focal severe stenosis in mid-left anterior descending coronary artery and a 60% to 70% stenosis in the proximal circumflex artery. Given advanced age and unstable hemodynamics, emergent multivessel PCI with complete revascularization was performed. An intra-aortic balloon pump was initially placed, but cardiogenic shock persisted despite this and multiple vasopressors. His femoral arteries were <5 mm bilaterally. Despite borderline access vessels, hemodynamic support with an Impella CP (Abiomed, Danvers, Massachusetts) was instituted via left common femoral access. The patient was only 4 feet 9 inches tall, weighed 50 kg with a body surface area of 1.4 m². Lack of flow distal to the common femoral arteriotomy was confirmed on duplex ultrasound exam (1). The patient was, however, completely dependent on the Impella CP and could not be weaned off despite maximal doses of 4 vasopressors (2,3).

It had already been established noninvasively and with a lower extremity angiogram (Figure 1) that the

patient had no antegrade distal flow beyond the site of insertion of the cannula, hence conservative management could threaten the viability of the limb. We thus decided to proceed with percutaneous temporary femoro-femoral bypass establishment by placing an antegrade sheath in the left superficial femoral artery and connecting it male-to-male with the contralateral common femoral sheath. This was

FIGURE 1 Lower Extremity Angiogram showing Occlusive Impella Cannula



The fluoro-masked "roadmap" image shows that the 13-F Impella CP cannula was occlusive for the common femoral artery of the diminutive index patient.

From the Division of Cardiology, Temple University Hospital, Philadelphia, Pennsylvania. Dr. Bashir is the cofounder of and has equity in Thrombolex Inc. Dr. O'Neill has received research support from Edwards Lifesciences. All other authors have reported that they have no relationships relevant to the contents of this paper to disclose.

Manuscript received February 28, 2017; revised manuscript received March 13, 2017, accepted March 23, 2017.

FIGURE 2 Successful Placement of Fully Percutaneous Extracorporeal Femoro-Femoral Bypass for Acute Limb Ischemia From Large-Bore Access



We were able to successfully establish flow distal to the insertion site of the occlusive arterial cannula by a temporary, novel percutaneous femoro-femoral bypass circuit.

achieved via careful antegrade access in the left superficial femoral artery, distal to the common femoral arteriotomy, and placement of a 6-F Brite Tip Cordis sheath (Cordis Corporation, Milpitas, California). This sheath was then connected male-to-male with the contralateral 6-F right common femoral artery retrograde sheath (Figure 2), thereby creating a temporary, novel percutaneous femoro-femoral bypass circuit (4). Subsequently, the left popliteal pulse was confirmed on physical exam, and duplex ultrasound suggested flow re-establishment in the left lower extremity.

ADDRESS FOR CORRESPONDENCE: Dr. Vikas Aggarwal, Section of Interventional Cardiology/ Division of Cardiology, Department of Internal Medicine, Lewis Katz School of Medicine at Temple University, Temple University Hospital, 3401 North Broad Street (C945), Philadelphia, Pennsylvania 19140. E-mail: Vikas.Aggarwal@tuhs.temple.edu.

REFERENCES

1. Abaunza M, Kabbani LS, Nypaver T, et al. Incidence and prognosis of vascular complications after percutaneous placement of left ventricular assist device. *J Vasc Surg* 2015;62:417-23.
2. Liu W, Mukku VK, Gilani S, Fujise K, Barbagelata A. Percutaneous hemodynamic support (Impella) in patients with advanced heart failure and/or cardiogenic shock not eligible to PROTECT II Trial. *Int J Angiol* 2013; 22:207-12.
3. Lima B, Kale P, Gonzalez-Stawinski GV, Kuiper JJ, Carey S, Hall SA. Effectiveness and safety of the Impella 5.0 as a bridge to cardiac transplantation or durable left ventricular assist device. *Am J Cardiol* 2016;117:1622-8.
4. Kizner L, Flottmann C, Horstkotte D, Gummert J. Bilateral antegrade perfusion of the superficial femoral artery to prevent limb ischaemia during combined use of Impella CP left ventricular assist device and extracorporeal life support. *Interact Cardiovasc Thorac Surg* 2016;23:335-7.

KEY WORDS acute limb ischemia, extracorporeal bypass, large bore access