

EDITORIAL COMMENT

## Frail Elderly, the Ideal Patients for MitraClip\*



Suzanne V. Arnold, MD, MHA

Frailty is a geriatric syndrome of impaired physiological reserve and decreased resistance to stressors (1). Characterized by a constellation of factors, including weakness, slowness, exhaustion, wasting, and low activity level, frailty is a risk factor for morbidity and mortality in a wide variety of situations independent of traditional demographic and clinical risk factors. From pneumonia (2) to myocardial infarction (3), general (4,5) and cardiac surgery (6,7), and even the less invasive procedure of transcatheter aortic valve replacement (8-10), a diagnosis of frailty increases the risk of mortality after nearly any intervention (11) or clinical insult. Furthermore, frailty increases the risk of failure to recover quality of life and functional status (10,12). An assessment of frailty has therefore been recommended for risk stratification before surgical intervention in elderly patients (10,11,13), but can be cumbersome and time-consuming, particularly when dealing with hospitalized patients. Although frailty assessments can be beneficial to manage patients' expectations for recovery, they are particularly important when they alter management. For example, an assessment may identify a patient as frail who would benefit from a less invasive approach to the treatment of their disease (e.g., multivessel coronary stenting or transcatheter aortic valve replacement instead of cardiac surgery). Or, it may identify patients who may benefit from pre-habilitation, early mobilization after intervention, or post-operative cardiac rehabilitation to

help the patient recover his or her functional status and preserve independence.

SEE PAGE 1920

In this issue of *JACC: Cardiovascular Interventions*, Metzke et al. (14) present their analysis where they performed a frailty assessment on 213 patients who were admitted for a MitraClip procedure at a single center in Germany. They found that frailty was present in nearly one-half of patients and also associated with an increased hazard of short- and long-term mortality, independent of demographic and clinical risk factors. These results were not surprising, because: 1) the MitraClip procedure is a procedure that is targeted at frail elderly patients who are not candidates for mitral valve surgery; and 2) frailty is associated with short- and long-term mortality in nearly every clinical scenario. The proportion of patients treated with a MitraClip who met criteria for frailty on the basis of standards set for community-dwelling elderly, although high, was perhaps not as high as expected. In patients with severe aortic stenosis and high surgical risk who were treated with transcatheter aortic valve replacement (a procedure that is more invasive than that of a MitraClip), 60% of patients met similar criteria for frailty (12). In these elderly patients who are felt to be at high surgical risk, the prevalence of frailty should be exceedingly high. Particularly in the case of mitral valve disease, patients who are not frail may, in fact, be better served by traditional cardiac surgery with more definitive treatment of the valve (at least in the case of degenerative mitral regurgitation). In this way, the assessment of frailty may help identify nonfrail patients who should have cardiac surgery instead of the MitraClip procedure.

Although the prevalence of frailty in this sample of patients treated with the MitraClip is not alarming (and may actually be on the low side), what is notable in this analysis is that frailty did not affect the

\*Editorials published in *JACC: Cardiovascular Interventions* reflect the views of the authors and do not necessarily represent the views of *JACC: Cardiovascular Interventions* or the American College of Cardiology.

From the Saint Luke's Mid America Heart Institute and University of Missouri-Kansas City, Kansas City, Missouri. Dr. Arnold is supported by a Career Development Grant Award (K23 HL116799) from the National Heart, Lung, and Blood Institute.

technical success nor the quality-of-life benefits of the procedure. In fact, frail patients were more likely than nonfrail patients to have a clinically relevant improvement in quality of life at 6 weeks after the MitraClip procedure. Much of this was driven by the fact that frail patients were more symptomatic and had substantially worse quality of life before the procedure and therefore had more to gain from mitral valve repair. However, it is reassuring that the procedure seemed to benefit the majority of patients, regardless of frailty status. For a procedure that has a primary goal to improve quality of life and is targeted at frail elderly patients, these findings support the continued use of MitraClip in this vulnerable population. Although the long-term survival of these frail patients is limited, palliative measures targeted at improvement in quality of life and functional status are still indicated.

The MitraClip procedure is a means to repair the mitral valve with little perioperative risk, but the results are suboptimal compared with traditional cardiac surgery (15). As such, this procedure should be limited

primarily to patients who are not candidates for more definitive treatment of their mitral regurgitation (16) (or, perhaps, in high-risk patients with functional mitral regurgitation). Frailty is a key reason to elect for a less invasive treatment option, and assessment of this may help determine which patients are ideal candidates for open mitral valve repair versus MitraClip. Results from Metzke et al. (14) suggest that although long-term survival is limited in frail patients, the MitraClip procedure can be done with technical success and can result in substantial improvement in quality of life. Although these data should be examined in a larger, multicenter study, they suggest that targeting the MitraClip procedure to frail patients can be an effective strategy to improve symptoms and quality of life with low perioperative risk.

---

**ADDRESS FOR CORRESPONDENCE:** Dr. Suzanne V. Arnold, Saint Luke's Mid America Heart Institute, 4401 Wornall Road, Kansas City, Missouri 64111. E-mail: [suz.v.arnold@gmail.com](mailto:suz.v.arnold@gmail.com).

---

## REFERENCES

1. Fried LP, Hadley EC, Walston JD, et al. From bedside to bench: research agenda for frailty. *Sci Aging Knowledge Environ* 2005;2005:pe24.
2. Torres OH, Munoz J, Ruiz D, et al. Outcome predictors of pneumonia in elderly patients: importance of functional assessment. *J Am Geriatr Soc* 2004;52:1603-9.
3. Ekerstad N, Swahn E, Janzon M, et al. Frailty is independently associated with short-term outcomes for elderly patients with non-ST-segment elevation myocardial infarction. *Circulation* 2011;124:2397-404.
4. Makary MA, Segev DL, Pronovost PJ, et al. Frailty as a predictor of surgical outcomes in older patients. *J Am Coll Surg* 2010;210:901-8.
5. Hewitt J, Moug SJ, Middleton M, et al. Prevalence of frailty and its association with mortality in general surgery. *Am J Surg* 2015;209:254-9.
6. Sundermann S, Dademasch A, Praetorius J, et al. Comprehensive assessment of frailty for elderly high-risk patients undergoing cardiac surgery. *Eur J Cardiothorac Surg* 2011;39:33-7.
7. Afialo J, Mottillo S, Eisenberg MJ, et al. Addition of frailty and disability to cardiac surgery risk scores identifies elderly patients at high risk of mortality or major morbidity. *Circ Cardiovasc Qual Outcomes* 2012;5:222-8.
8. Stortecky S, Schoenenberger AW, Moser A, et al. Evaluation of multidimensional geriatric assessment as a predictor of mortality and cardiovascular events after transcatheter aortic valve implantation. *J Am Coll Cardiol Intv* 2012;5:489-96.
9. Schoenenberger AW, Stortecky S, Neumann S, et al. Predictors of functional decline in elderly patients undergoing transcatheter aortic valve implantation (TAVI). *Eur Heart J* 2013;34:684-92.
10. Green P, Arnold SV, Cohen DJ, et al. Relation of frailty to outcomes after transcatheter aortic valve replacement (from the PARTNER trial). *Am J Cardiol* 2015;116:264-9.
11. Lin HS, Watts JN, Peel NM, Hubbard RE. Frailty and post-operative outcomes in older surgical patients: a systematic review. *BMC Geriatr* 2016;16:157.
12. Arnold SV, Afialo J, Spertus JA, et al., U.S. CoreValve Investigators. Prediction of poor outcome after transcatheter aortic valve replacement. *J Am Coll Cardiol* 2016;68:1868-77.
13. Afialo J, Kim S, O'Brien S, et al. Gait speed and operative mortality in older adults following cardiac surgery. *JAMA Cardiol* 2016;1:314-21.
14. Metzke C, Matzik A-S, Scherner M, et al. Impact of frailty on outcomes in patients undergoing percutaneous mitral valve repair. *J Am Coll Cardiol Intv* 2017;10:1920-9.
15. Mauri L, Foster E, Glower DD, et al., EVEREST II Investigators. 4-year results of a randomized controlled trial of percutaneous repair versus surgery for mitral regurgitation. *J Am Coll Cardiol* 2013;62:317-28.
16. Nishimura RA, Otto CM, Bonow RO, et al. 2017 AHA/ACC focused update of the 2014 AHA/ACC guideline for the management of patients with valvular heart disease: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol* 2017;70:252-89.

---

**KEY WORDS** frailty, mitral valve disease, quality of life