

CORRECTIONS

Räber L, Baumgartner S, Garcia Garcia HM, Kalesan B, Justiz J, Pilgrim T, Moschovitis A, Khattab AA, Buellesfeld L, Wenaweser P, Meier B, Serruys PW, Jüni P, Windecker S. Long-Term Vascular Healing in Response to Sirolimus- and Paclitaxel-Eluting Stents: An Optical Coherence Tomography Study. *J Am Coll Cardiol Interv* 2012;5:946–57.

Please note that Dr. Garcia Garcia’s name was misspelled in the original article.

The correct version should be:
 Hector M. Garcia-Garcia.

The authors apologize for this error.

<http://dx.doi.org/10.1016/j.jcin.2012.10.001>

Rao SV, Dai D, Subherwal S, Weintraub WS, Brindis RS, Messenger JC, Lopes RD, Peterson ED. Association Between Periprocedural Bleeding and Long-Term Outcomes Following Percutaneous Coronary Intervention in Older Patients. *J Am Coll Cardiol Interv* 2012;5:958–65.

Figure 2 contained an error in the legend and in the explanatory text making the figure difficult to interpret. The corrected version of the legend and explanatory text is shown here:




Figure 2. Kaplan-Meier Rates

(A) Kaplan-Meier rates of readmission for bleeding between patients who developed periprocedural bleeding and those who did not. (B) Kaplan-Meier rates of major adverse cardiac events (MACE) between patients who developed periprocedural bleeding and those who did not. (C) Kaplan-Meier rates of all-cause mortality between patients who developed periprocedural bleeding and those who did not.