

IMAGES IN INTERVENTION

Percutaneous Treatment of Chronic Distal Aortic Occlusion

A Viable Option




Mohamad Alkhouli, MD, Irfan Shafi, MD, Pravin Patil, MD, Riyaz Bashir, MB BS

A 55-year-old woman presented with severe lifestyle-limiting bilateral calf and thigh claudication (Rutherford stage III). She had history of aortobifemoral bypass 13 years ago, which had occluded since then. Her ankle brachial index (ABI) was 0.67 on the right and 0.69 on the left. Computed tomography (CT) and digital-subtraction angiograms showed an occluded infrarenal aorta (**Figures 1A and 2A, Online Video 1**). She was deemed

to be a poor candidate for surgical revascularization secondary to her comorbidities (uncontrolled diabetes and severe coronary artery disease) and the need for redo surgery.

She was treated successfully with percutaneous bilateral kissing aortoiliac stenting using bilateral femoral and left radial approaches. The left aortoiliac occlusion was successfully traversed in a retrograde fashion from the left femoral access using a Pioneer






FIGURE 2 Aortoiliac Angiogram

(A) Occluded distal aorta. (B) Reconstruction of the aortoiliac bifurcation with kissing stents.

re-entry catheter (Volcano, San Diego, California). The right aortoiliac occlusion was traversed in an antegrade fashion from the left radial approach using an 0.018-inch Victory wire (Boston Scientific, Natick, Massachusetts) that was externalized via the right femoral sheath. The aortoiliac bifurcation was reconstructed with kissing balloon angioplasty and stenting with balloon-expandable Assurant-Cobalt stents (Medtronic, Langhorne, Pennsylvania) and self-expanding LifeStar stents (Bard Peripheral Vascular, Tempe, Arizona) with excellent angiographic and hemodynamic results (Figures 1B and 2B, Online Video 2). The following day, the patient was walking without any claudication and has continued to be free of symptoms at 3-month follow-up. Repeat

ABIs were completely normal (1.17 on the left and 1.19 on the right side).

Surgical revascularization is considered to be the preferred treatment of infrarenal aortic occlusion (TASC-D lesion) (1). However, with current advances in percutaneous technologies such as intravascular ultrasound-guided re-entry devices, endovascular treatment seems to be a very reasonable first line approach particularly in high-risk surgical patients (2,3).

REPRINT REQUESTS AND CORRESPONDENCE: Dr. Mohamad Alkhouli, Division of Cardiovascular Diseases, Temple University School of Medicine, 3401 North Broad Street (9 PP), Philadelphia, Pennsylvania 19140. E-mail: adnanalkhouli@gmail.com.

REFERENCES

1. Norgren L, Hiatt WR, Dormandy JA, et al. Inter-Society Consensus for the Management of Peripheral Arterial Disease (TASC II). *Eur J Vasc Endovasc Surg* 2007;33 Suppl 1:S1-75.
2. Moise MA, Alvarez-Tostado JA, Clair DG, et al. Endovascular management of chronic infrarenal aortic occlusion. *J Endovasc Ther* 2009;16:84-92.
3. Kim TH, Ko YG, Kim U, et al. Outcomes of endovascular treatment of chronic total occlusion of the infrarenal aorta. *J Vasc Surg* 2011;53:1542-9.

KEY WORDS chronic total occlusion, distal aortic occlusion, endovascular, re-entry devices

APPENDIX For supplemental videos and their legends, please see the online version of this article.