

CRT-203

Safety and Clinical Outcomes of Endovascular Treatment for Extrinsic Pulmonary Artery Stenosis: A Systematic Review

Mohammed A. Chowdhury, Mujeeb A. Sheikh, George V. Moukarbel
University of Toledo, Toledo, OH

Background: Endovascular stent implantation for extrinsic pulmonary arterial stenosis (PAS) in adults is anecdotal.

Method: We systematically reviewed etiology, clinical characteristics, safety and outcomes of the published cases of extrinsic PAS treated with endovascular approach.

Result: A total of 113 published cases of extrinsic PAS were reviewed. The most common etiology of extrinsic PAS was mediastinal mass (31%), primary tumors of pulmonary artery (27%), fibrosing mediastinitis (18%), pulmonary pseudoaneurysm (17%), anastomotic stenosis post lung transplantation (7%) and radiation induced pulmonary stenosis (1%). Mean age at presentation was 52 years \pm 14.75 with a M:F ratio of 1:1. Dyspnea (41%) was the most common presenting symptom followed by chest pain (29%), hemoptysis (11%), cough (11%), peripheral edema (7.4%), right heart failure (6%) and pulmonary hypertension (3%). Right Pulmonary artery was involved in 40%, bilateral pulmonary artery in 35% and left pulmonary artery in 15% of cases. 41(36%) cases reported stent implantation with Palmaz, Wall stent and stent grafts being commonly used stents. In all cases endovascular treatment was associated with symptom improvement along with reduction in trans-lesional gradient. 6 (15%) cases reported immediate complications after stent implantation. In 7 (17%) cases in-stent restenosis was reported at 6 months follow-up. There was no immediate periprocedural mortality.

Conclusion: Stent implantation is an effective and safe alternative option for the treatment of extrinsic PAS.

CRT-204

Same Day Discharge Right and Left Catheterization to Test Vasoreactivity in Pulmonary Hypertension by Basilar and Radial Access

Luis Enrique Berumen Dominguez, Jose Luis Ojeda-Delgado, Hugo Gutierrez-Leonard,
Luis Manuel Paez-Lizarraga, Miguel Ramirez-Aldaraca
Hospital Militar, Mexico, Mexico

Background: Pulmonary Hypertension is a frequent disease in third level hospitals, with elevated morbidity and mortality in short time. To choose the medical treatment is fundamental to do a pulmonary vasoreactivity test.

Objective: To show and classify the results of the vasoreactivity test in the different groups of pulmonary Hypertension, done by basilar and radial Access.

Material and Method: By consecutive simple we captured cases in which left and right catheterism is done in patients with pulmonary hypertension, since November 2010 to September 2013. By basilar access we did the right catheterism, oximetries and pressure are registered, in superior vena cava, Inferior vena cava, Right Ventricle and Pulmonary Trunk. By radial Access pressures and oximetries are registered in aorta and Left Ventricle. Cardiac output is determined by Fick Method. The vasoreactivity was evaluated with adenosine infusion in the pulmonary trunk, in doses of 100 mcg/kg/min with a maximum of 12 mgs. A positive test is considered when Mean Pulmonary arterial pressure (mPAP) decrease more than 10 mmHg or a (mPAP) less than 40 mmHg without affection of the cardiac output.

Results: 114 procedures were done, 57% women, mean age 44 years, 74 patients of the group 1, 26 of the group 2, 8 of the group 3 and 6 of the group 4. Only 12% of the patients responded to adenosine infusion, 11 patients from the group 1 and 3 patients from de group 3 (Pulmonary Hypertension classification, Dana Point 2008). The mPAP decreased on average of 17 mmHg, and it was sustained in average 7.2 minutes. All procedures were done successfully and same day discharge. Complications: 2 patients referred severe chest pain with the adenosine infusion, but not persistent electrocardiographic abnormalities are founded. 10% of patients presented pain in the Access site that was controlled with analgesics.

Conclusions: Basilar and radial Access was accepted in 100% percent of the patients, and all are same day discharged. There were no major complications and the vasoreactivity test was positive only in 8% of the patients, principally patients of the group 1.

Neurovascular Intervention

CRT-205

Impact of Veins Endovascular Procedures on the Quality of Life in Patients with Multiple Sclerosis

Marjeta Zorc,¹ Miro Denislić,² Zoran Milosevic,³ Rok Zorec,¹ Dean Ravnik,¹
Ruda Zorc Pleskovic,¹ Ales Pleskovic,³ Oscar Mendiz⁴

¹University of Ljubljana, Medical Faculty, Ljubljana, Slovenia; ²Medical Centre Medicor, Ljubljana, Slovenia; ³University Clinical Centre Ljubljana, Ljubljana, Slovenia;

⁴Favaloro foundation, Buenos Aires, Argentina

Background: Chronic cerebrospinal venous insufficiency (CCSVI) is a new challenge of extracranial venous pathway, which provides at present still controversial insight into the vascular role of multiple sclerosis (MS). The aim of our open-label study was to evaluate quality of life (QoL) of MS patients after endovascular procedures.

Materials and Methods: MS patients diagnosed by revised McDonald criteria who fulfilled Doppler sonography criteria for CCSVI. To evaluate the efficacy of the vascular procedures on QoL Multiple Sclerosis Impact Scale (MSIS-29), Fatigue Severity Scale (FSS), Modified Fatigue Impact Scale (MFIS) and Overactive bladder self-administered questionnaire (OAB-V8) were used. MSIS-29, FSS, MFIS and OAB-V8 were evaluated at baseline, 3, 6, and 12 months after vascular angioplasty.

Results: In our study 72 consecutive MS patients were included. FSS ($p < 0.001$) and MFIS scores - total score, as well as three subscale scores (physical, psychosocial and cognitive) significantly improved after vascular procedures ($p < 0.05$). The physical subscale correlated with the degree of pyramidal impairment. The important improvement of the bladder function using OAB-V8 ($p < 0.01$) and QoL assessed by MSIS-29 questionnaire ($p < 0.01$) were obtained.

Conclusions: The endovascular procedure demonstrated a beneficial effect on the quality of life of MS patients. The amelioration of cerebral venous drainage significantly reduced the perception of fatigue, increased the mental health and emotional stability, respectively. Additionally, important improvement of the bladder dysfunction even in MS patients with the progressive course of disease was achieved. Also, the sexual function improved. The relationship between CCSVI and hypoperfusion was demonstrated. The role of better cerebrospinal fluid flow after vascular procedures is discussed. It seems that the endovascular procedures in MS patients may influence the clinical picture of MS patients. Further investigations also in other neurodegenerative diseases are recommended.

CRT-206

Safety and Efficacy of Carotid Stenting in the Treatment of Carotid Artery Stenosis: Immediate Results and Long Term Follow-Up in Our Experience

Cinzia Moncalvo,¹ Vincenzo Puma,¹ Mario Caprile,¹ Angelo Laurenza,¹
Giuseppe Celano,² Eugenio Novelli,² Giuseppe Carosio,¹ Paolo Cioffi¹

¹NCC Città di Alessandria, Alessandria, Italy; ²Clinica San Gaudenzio, Novara, Italy

Background: Carotid artery stenting (CAS) may be an alternative to surgical endarterectomy for the treatment of atherosclerotic carotid artery stenosis.

Purpose: to analyze retrospectively the procedures of CAS performed in our Centre between January 2004 and December 2012.

Methods: This analysis includes 604 procedures performed in 554 patients (382 men; mean age: 72 years old). Symptomatic patients with carotid artery stenosis $> 50\%$ were the 45%; we treated asymptomatic patients affected by $> 70\%$ stenosis. 398 patients (72%) were considered at high surgical risk: 207 (37%) with severe contralateral stenosis; 136 (24%) with severe or unstable angina, poor left ventricular function, left main disease or trivascular coronary artery disease, severe cardiac valve disease; 35 patients (6%) presented restenosis after surgical treatment. 28 (5%) patients were