

## EDITORIAL COMMENT

# The Rise and Fall of Aspiration Thrombectomy\*



Dharam J. Kumbhani, MD, SM,<sup>†</sup> Anthony A. Bavry, MD, MPH<sup>‡</sup>

The hallmark of ST-segment elevation myocardial infarction (STEMI) is thrombotic occlusion of an epicardial vessel. As many as 50% of patients undergoing primary percutaneous coronary intervention (PCI) demonstrate distal embolization and microvascular obstruction despite achieving normalized epicardial flow. This is associated with increased infarct size, reduced recovery of ventricular function, and reduced survival (1,2). In an effort to reduce distal embolization, a number of thrombectomy devices have been studied for use with primary PCI. The story of aspiration thrombectomy is a sterling example of the utility of evidence-based medicine in contemporary clinical practice, where initial exuberance for a new therapy was tempered by a more limited or measured application on the basis of accrual of high-quality data. Some of the salient features are discussed in the following text.

## THE RISE

TAPAS (Thrombus Aspiration during Percutaneous Coronary Intervention in Acute Myocardial Infarction Study) was one of the sentinel trials in this field (n = 1,071). At 1 year, there were significant reductions in all-cause (4.7% vs. 7.6%; p = 0.04) and cardiovascular (3.6% vs. 6.7%; p = 0.02) mortality, with a trend toward reduction in subacute and late stent thrombosis with adjunctive aspiration thrombectomy compared with conventional PCI

\*Editorials published in *JACC: Cardiovascular Interventions* reflect the views of the authors and do not necessarily represent the views of *JACC: Cardiovascular Interventions* or the American College of Cardiology.

From the <sup>†</sup>Division of Cardiology, University of Texas Southwestern Medical Center, Dallas, Texas; and the <sup>‡</sup>Division of Cardiology, University of Florida, Gainesville, Florida. Both authors have received honoraria from the American College of Cardiology.

alone (3). Although the trial was criticized for its limitations (possible selection bias [single-center study], unclear mechanism [no difference in enzymatic infarct size post-procedure but a significant difference in cardiovascular and all-cause mortality at 1 year], and possibility of chance [trial was not powered for mortality]), the overall clinical outcomes were directionally consistent with other smaller trials (2). On the basis of this data, aspiration thrombectomy received a Class IIa recommendation for use with PCI in the 2009 American College of Cardiology (ACC)/ American Heart Association (AHA) STEMI guidelines (4).

SEE PAGE 126

Commensurate with the timeframe of evidence accrual, there appeared to be a gradual uptake of aspiration thrombectomy in clinical practice the world over. In this issue of *JACC: Cardiovascular Interventions*, Sirker et al. (5) estimate that the uptake of adjunctive thrombectomy in the United Kingdom was around 50% by 2010 to 2011.

## THE FALL

The INFUSE-AMI (Infuse-Acute Myocardial Infarction) trial failed to replicate the findings of the TAPAS trial in 452 patients, with no differences observed in infarct size with aspiration thrombectomy at 30 days (6). Despite this, on systematic review, the totality of evidence still favored a benefit in clinical outcomes with aspiration thrombectomy, particularly on longer-term follow-up (7), and it remained a Class IIa recommendation in the 2013 ACC/AHA STEMI guidelines (8).

TASTE (Thrombus Aspiration in ST-Elevation Myocardial Infarction in Scandinavia) was a large, multicenter, registry-based randomized clinical trial (n = 7,244). There were no differences in clinical

outcomes assessed at 1 year between aspiration thrombectomy versus conventional PCI, including all-cause mortality (5.3% vs. 5.6%;  $p = 0.57$ ) or stent thrombosis (0.7% vs. 0.9%;  $p = 0.51$ ) (9). TOTAL (Trial of Routine Aspiration Thrombectomy with PCI versus PCI Alone in Patients with STEMI) is the largest and most definitive trial on this topic to date ( $n = 10,732$ ). No differences in clinical outcomes were observed at 6 months in favor of aspiration thrombectomy, including in the primary outcome of major adverse cardiovascular events (6.9% vs. 7.0%;  $p = 0.86$ ), cardiovascular death (3.1% vs. 3.5%;  $p = 0.34$ ), or stent thrombosis (1.5% vs. 1.7%;  $p = 0.42$ ); stroke rates were higher with aspiration thrombectomy (0.7% vs. 0.3%;  $p = 0.02$ ) (10). On the basis of this evidence, routine aspiration thrombectomy has been downgraded to a Class III recommendation in the most recent (2015) iteration of the ACC/AHA STEMI guidelines (11).

The current study by Sirker et al. (5) provides further evidence against routine aspiration thrombectomy use during primary PCI. They included 98,176 STEMI patients undergoing primary PCI from the U.K. national PCI registry. On propensity analysis, there was no difference in 1-year mortality between any form of thrombectomy versus conventional PCI, including adjunctive aspiration thrombectomy and conventional PCI ( $n = 85,675$ ;  $p = 0.96$ ) (5). This is an important body of work on a large cohort of patients. However, causative inferences regarding therapeutic choices from observational datasets can be hard at best, and outright wrong at worst, as unmeasured confounders can obfuscate all and any statistical adjustments. Physicians frequently make therapeutic decisions on the basis of characteristics that can be hard to capture in registries (such as frailty, “eyeball test,” and so on). In such settings, inverse probability of treatment weights may be a less confounded way of performing a propensity analysis, wherein individuals are weighted by the inverse probability

of receiving the treatment that they actually received. In the current study, the authors chose to perform a more conventional propensity analysis, thus being potentially more vulnerable to selection biases. However, the results of the current study are directionally consistent with recent randomized controlled trials, and are thus less likely to be spurious. Coding errors may also be present in registries. For example, patients in this study that did not receive thrombectomy were older, more frequently had diabetes, less frequently smoked, and more frequently had pre-procedure Thrombolysis In Myocardial Infarction 2 to 3 flow. Although these patients were classified as “STEMI,” such characteristics more closely resemble that of non-STEMI, which carries a different prognosis than STEMI. Left ventricular ejection fraction was also missing for a large proportion of patients, which is an important determinant of outcomes following STEMI, and may also result in unmeasured confounding. However, the authors performed sensitivity analyses using both nonimputed and imputed data for LVEF and report a similar treatment effect (5).

In summary, routine aspiration thrombectomy was once considered a relatively simple technique to improve long-term outcomes in the care of STEMI patients. Thrombus removal remains an intuitively valid concept; however, thrombus removal is more complex than initially appreciated. Future studies will need to similarly rigorously evaluate the role of selective thrombectomy versus no thrombectomy, determine more effective thrombectomy devices, and better discriminate appropriate lesions to which this therapy can be applied.

**REPRINT REQUESTS AND CORRESPONDENCE:** Dr. Dharam J. Kumbhani, UT Southwestern Medical Center, 5323 Harry Hines Boulevard, Dallas, Texas 75390-9047. E-mail: [dharam@post.harvard.edu](mailto:dharam@post.harvard.edu).

## REFERENCES

1. Poli A, Fetiveau R, Vandoni P, et al. Integrated analysis of myocardial blush and ST-segment elevation recovery after successful primary angioplasty: real-time grading of microvascular reperfusion and prediction of early and late recovery of left ventricular function. *Circulation* 2002;106:313-8.
2. Bavry AA, Kumbhani DJ, Bhatt DL. Role of adjunctive thrombectomy and embolic protection devices in acute myocardial infarction: a comprehensive meta-analysis of randomized trials. *Eur Heart J* 2008;29:2989-3001.
3. Vlaar PJ, Svilaas T, van der Horst IC, et al. Cardiac death and reinfarction after 1 year in the Thrombus Aspiration during Percutaneous coronary intervention in Acute myocardial infarction Study (TAPAS): a 1-year follow-up study. *Lancet* 2008;371:1915-20.
4. Kushner FG, Hand M, Smith SC Jr., et al. 2009 focused updates: ACC/AHA guidelines for the management of patients with ST-elevation myocardial infarction (updating the 2004 guideline and 2007 focused update) and ACC/AHA/SCAI guidelines on percutaneous coronary intervention (updating the 2005 guideline and 2007 focused update) a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* 2009;54:2205-41.
5. Sirker A, Mamas M, Kwok CS, Kontopantelis E, Ludman P, Hildick-Smith D, on behalf of BCIS. Outcomes from selective use of thrombectomy in patients undergoing primary percutaneous coronary intervention for ST-segment elevation myocardial infarction: an analysis of the British Cardiovascular Intervention Society/National Institute for Cardiovascular Outcomes Research (BCIS-NICOR) Registry 2006-2013. *J Am Coll Cardiol Intv* 2016;9:126-34.
6. Stone GW, Maehara A, Witzenbichler B, et al. Intracoronary abciximab and aspiration thrombectomy in patients with large anterior myocardial infarction: the INFUSE-AMI randomized trial. *JAMA* 2012;307:1817-26.

7. Kumbhani DJ, Bavry AA, Desai MY, Bangalore S, Bhatt DL. Role of aspiration and mechanical thrombectomy in patients with acute myocardial infarction undergoing primary angioplasty: an updated meta-analysis of randomized trials. *J Am Coll Cardiol* 2013;62:1409-18.
  8. O'Gara PT, Kushner FG, Ascheim DD, et al. 2013 ACCF/AHA guideline for the management of ST-elevation myocardial infarction: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* 2013;61:e78-140.
  9. Lagerqvist B, Frobert O, Olivecrona GK, et al. Outcomes 1 year after thrombus aspiration for myocardial infarction. *N Engl J Med* 2014;371:1111-20.
  10. Jolly SS, Cairns JA, Dzavik V. Primary PCI with or without thrombectomy. *N Engl J Med* 2015;373:682-3.
  11. Levine GN, O'Gara PT, Bates ER, et al. 2015 ACC/AHA/SCAI focused update on primary percutaneous coronary intervention for patients with ST-elevation myocardial infarction: an update of the 2011 ACCF/AHA/SCAI Guideline for Percutaneous Coronary Intervention and the 2013 ACCF/AHA Guideline for the Management of ST-Elevation Myocardial Infarction: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Society for Cardiovascular Angiography and Interventions. *J Am Coll Cardiol* 2015 Oct 21 [E-pub ahead of print].
- 
- KEY WORDS** myocardial infarction, outcomes, registry, thrombectomy