

EDITORIAL COMMENT

# Thoracic Endovascular Aortic Repair in Acute and Chronic Type B Aortic Dissection\*



Santi Trimarchi, MD, PhD,<sup>†</sup> Kim A. Eagle, MD<sup>‡</sup>

The temporal course of type B aortic dissection (TBAD) can be divided into several phases: hyperacute (<24 h), acute (within 2 weeks), subacute (2 weeks up to 3 months), and chronic (>3 months) (1-3). This distinction has become highly important, because it may help to identify the time interval during which these patients are at higher risk for complications and, for selected patients, the optimal moment for TEVAR (Thoracic Endovascular Aortic Repair). Identification of both the optimal cohort and most ideal time to perform TEVAR remains a matter of debate.

The VIRTUE trial suggested that ideal timing of TEVAR should be in the subacute phase, when the aorta still has aortic plasticity, which leads to similar remodeling and survival rates compared with treatment in the acute setting (3). In addition, it may also be that the incidence of TEVAR-related complications, like retrograde dissection, is reduced by waiting a period of weeks after acute type B dissection.

In chronic TBAD, anatomic findings, such as narrow lumens and thickening of the intimal flap, may inhibit optimal aortic remodeling after TEVAR (4). Although open repair has higher operative mortality and morbidity, the previously noted anatomic issues are not insurmountable. However, no definitive evidence currently exists to confirm which method is optimal in uncomplicated type B dissection.

Currently, the optimal management for acute complicated TBAD is felt to be TEVAR, whereas open surgery is reserved for younger patients, patients with connective tissue disease such as Marfan syndrome, and those unsuitable for endovascular repair. For uncomplicated type B dissection, there is an ongoing debate regarding the most suitable therapy: medical therapy alone or endovascular along with optimal medical treatment. Although the ADSORB (Acute Dissection treatment with Stent graft OR Best medical therapy) and INSTEAD (the INvestigation of STEnt grafts in Aortic Dissection) trials showed long-term benefits of TEVAR in terms of survival and remodeling (5,6), several critiques have been published about these trials, mainly regarding small sample size and the primary outcome measures. Because of this, a conservative approach for many patients with uncomplicated TBAD is still encouraged.

In chronic TBAD, the indication for treatment is usually aneurysmal degeneration of the dissected segments of the aorta after initial medical therapy, which extends into the abdominal of the aorta in almost two-thirds of these patients (7-9).

After standard TEVAR, which includes stent graft treatment limited to the descending aorta, complete false lumen thrombosis during follow-up is described between 39% and 100% of patients (10), and rates of favorable aortic remodeling range between 5% and 89% (11-13). In these patients, despite remodeling in the proximal stented sections, distal segments of the aorta may still be patent.

SEE PAGE 183

\*Editorials published in *JACC: Cardiovascular Interventions* reflect the views of the authors and do not necessarily represent the views of *JACC: Cardiovascular Interventions* or the American College of Cardiology.

From the <sup>†</sup>University of Milan, Policlinico San Donato I.R.C.C.S., Cardiovascular Department, Thoracic Aortic Research Center, San Donato Milanese, Italy; and the <sup>‡</sup>University of Michigan Health System, Cardiovascular Center, Ann Arbor, Michigan. Dr. Trimarchi has served as a consultant and speaker for Medtronic and W.L. Gore. Dr. Eagle has received research grants from W.L. Gore, Medtronic, and Terumo.

In the paper by Fanelli et al. (14) in this issue of *JACC: Cardiovascular Interventions*, predictors and influencing factors of remodeling and outcome after TEVAR for both acute and chronic TBAD are

addressed. They used the “old” definition of acute and chronic ( $\leq 14$  or  $>14$  days after onset). The results could have been different if they had used truly chronic patients, namely after 3 months from the onset of symptoms. The factors they assessed are interesting, because some of these have not been studied well before, such as oversizing, the stent, left subclavian artery embolization, and length and number of implanted devices. Interestingly, oversizing did not influence the occurrence of endoleaks, remodeling, or major adverse events, although 2 cases of retrograde dissection were reported in the acute cohort. Furthermore, the length of coverage did not have any influence on remodeling. In chronic dissections, it may be expected that the length of coverage would be of importance in type IIIb patients, because a patent abdominal aortic segment may lead to aortic growth and inhibition of favorable aortic remodeling. In the current results, however, endoleaks were not related to the degree of false lumen thrombosis in the chronic patients.

A significant enlargement of the aortic diameter above the stent graft was seen in the acute group, which is important because this may reflect increased strain on the aorta in the segments adjacent to the stent graft. This effect of the stent graft on the aorta and its hemodynamic surroundings should be studied in more detail.

The observation that left subclavian artery embolization was protective against endoleak development in acute cases is interesting, because it promoted false lumen thrombosis. This was not observed in the chronic TBAD cohort.

Of course, these results need to be interpreted carefully, because it is a retrospective case series with a limited number of patients, including 5 cases of post-traumatic dissection that per se are thought to have a different etiology, prognosis, and evolution than spontaneous dissection. It seems pretty clear that precise characterization of the temporal events, including the exact time between the onset and the intervention, is critical. It appears that TEVAR in a 2-day-old and 3-week-old dissection can give significantly different results compared with a 12-month-old dissection in terms of stent graft-related risks, aortic remodeling, and technical success. The focus of the current paper is on diameter only; however, a volumetric analysis of the true and false lumen and the size and precise location of the entry tear would have added even more value to the paper, because this gives a more clear indication of the changes in both lumens after TEVAR.

Nevertheless, given the paucity of studies looking at this cohort, this paper adds to the knowledge and understanding of the differences between TEVAR for acute and chronic type B aortic dissections in terms of management and remodeling.

Usefulness of an uncovered stent graft in the abdominal aorta following standard TEVAR has become a viable option for treating complicated acute TBAD with malperfusion (15). Mid-term outcomes in this group of patients have been associated with promising results in terms of aortic remodeling; however, no randomized and controlled studies have been reported on its routine use.

Branched and fenestrated TEVAR has also become an emerging technique to treat patients with chronic TBAD with less favorable anatomy with endovascular management (16). These devices, which are patient-specific with dedicated fenestrations and branches, provide an opportunity to treat a larger cohort of patients with less invasive tools. Early reports surrounding branched and fenestrated TEVAR seem promising, but long-term results in the thoracic aorta are limited. Studies of larger cohorts with longer follow-up are widely anticipated. Currently, these devices can only be used in an elective setting, because the patient-specific fabrication takes time. In the future, it is conceivable that off-the-shelf branched and fenestrated devices will become available to treat emergent cases as well.

In addition to off-the-shelf devices, development of branched and fenestrated stent grafts specifically for chronic TBADs are needed, because continued perfusion of the distal aortic segments is often described, and stenting a longer section of the descending aorta may be necessary. This represents a problem with the current stent grafts, because they are not designed to also cover the visceral segments.

The entire field of TEVAR for aortic dissection is rapidly evolving. We need better technology, more randomized trials, and more precise phenotyping as we strive to optimize patient outcomes.

---

**REPRINT REQUESTS AND CORRESPONDENCE:** Dr. Santi Trimarchi, University of Milan, Thoracic Aortic Research Center, Vascular Surgery Unit, Cardiovascular Department, Policlinico San Donato I.R.C.C.S., Piazza Malan 2, 20097, San Donato Milanese, Italy. E-mail: [santi.trimarchi@unimi.it](mailto:santi.trimarchi@unimi.it) OR [santi.trimarchi@grupposandonato.it](mailto:santi.trimarchi@grupposandonato.it).

---

**REFERENCES**

1. Booher AM, Isselbacher EM, Nienaber CA, et al. The IRAD classification system for characterizing survival after aortic dissection. *Am J Med* 2013;126:730.e19-24.
2. Dake MD, Thompson M, van Sambeek M, et al. DISSECT: a new mnemonic-based approach to the categorization of aortic dissection. *Eur J Vasc Endovasc Surg* 2013;46:175-90.
3. VIRTUE Registry Investigators. Mid-term outcomes and aortic remodelling after thoracic endovascular repair for acute, subacute, and chronic aortic dissection: the VIRTUE Registry. *Eur J Vasc Endovasc Surg* 2014;48:363-71.
4. Erbel R, Aboyans V, Boileau C, et al., for the Task Force for the Diagnosis and Treatment of Aortic Diseases of the European Society of Cardiology (ESC). 2014 ESC guidelines on the diagnosis and treatment of aortic diseases: document covering acute and chronic aortic diseases of the thoracic and abdominal aorta of the adult. *Eur Heart J* 2014;35:2873-926.
5. Brunkwall J, Kasprzak P, Verhoeven E, et al. Endovascular repair of acute uncomplicated aortic type b dissection promotes aortic remodelling: 1 year results of the ADSORB Trial. *Eur J Vasc Endovasc Surg* 2014;48:205-91.
6. Nienaber CA, Kische S, Rousseau H, et al. Endovascular repair of type B aortic dissection: long-term results of the randomized investigation of stent grafts in aortic dissection trial. *Circ Cardiovasc Interv* 2013;6:407-16.
7. Guangqi C, Xiaoxi L, Wei C, et al. Endovascular repair of Stanford type B aortic dissection: early and mid-term outcomes of 121 cases. *Eur J Vasc Endovasc Surg* 2009;38:422-6.
8. Lee M, Lee do Y, Kim MD, et al. Outcomes of endovascular management for complicated chronic type B aortic dissection: effect of the extent of stent graft coverage and anatomic properties of aortic dissection. *J Vasc Interv Radiol* 2013;24:1451-60.
9. Andacheh ID, Donayre C, Othman F, et al. Patient outcomes and thoracic aortic volume and morphologic changes following thoracic endovascular aortic repair in patients with complicated chronic type B aortic dissection. *J Vasc Surg* 2012;56:644-50, discussion 50.
10. Thrumurthy SG, Karthikesalingam A, Patterson BO, et al. A systematic review of mid-term outcomes of thoracic endovascular repair (TEVAR) of chronic type B aortic dissection. *Eur J Vasc Endovasc Surg* 2011;42:632-47.
11. Jia X, Guo W, Li TX, et al. The results of stent graft versus medication therapy for chronic type B dissection. *J Vasc Surg* 2013;57:406-14.
12. Oberhuber A, Winkle P, Schelzig H, et al. Technical and clinical success after endovascular therapy for chronic type B aortic dissections. *J Vasc Surg* 2011;54:1303-9.
13. van Bogerijen GH, Patel HJ, Williams DM, et al. Propensity adjusted analysis of open and endovascular thoracic aortic repair for chronic type B dissection: a twenty-year evaluation. *Ann Thorac Surg* 2015;99:1260-6.
14. Fanelli F, Cannavale A, O'Sullivan GJ, et al. Endovascular repair of acute and chronic aortic Type B dissections: main factors affecting aortic remodeling and clinical outcome. *J Am Coll Cardiol Intv* 2016;9:183-91.
15. Lombardi JV, Cambria RP, Nienaber CA, et al. Aortic remodeling after endovascular treatment of complicated type B aortic dissection with the use of a composite device design. *J Vasc Surg* 2014;59:1544-54.
16. Oikonomou K, Kopp R, Katsargyris A, et al. Outcomes of fenestrated/branched endografting in post-dissection thoracoabdominal aortic aneurysms. *Eur J Vasc Endovasc Surg* 2014;48:641-8.

---

**KEY WORDS** aortic dissection, TEVAR, type B dissection